

ArmorTech®

GILLOTINE™

For foliar and systemic control of insects in commercial and residential landscapes and interiorscapes, non-bearing fruit and nut trees, residential apple and pear trees, turfgrass and other sites where plants are grown for ornamental, aesthetic and climate modification purposes

Active Ingredient	By Wt.
* Clothianidin	50.0%
Other Ingredients	50.0%
Total	100.0%
*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine	

EPA Reg. No. 59639-152-86064
 EPA Est. 39578-TX-01 (C)
 67545-AZ-01 (V) Superscript
 is first letter in lot number.

**KEEP OUT OF
 REACH OF
 CHILDREN
 CAUTION**

SEE INSIDE BOOKLET FOR
 ADDITIONAL PRECAUTIONARY
 STATEMENTS.

**NET WEIGHT
 2 LB 8 OZ**

FIRST AID

- If in eyes:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If swallowed:**
- Call poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If on skin or clothing:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If inhaled:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

Manufactured for **United Turf Alliance, LLC**
 8014 Cumming Highway, Suite 403-282
 Canton, GA 30115

Form 1787-A



United Turf Alliance

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

FIRST AID

- | | |
|--------------------------------|---|
| If in eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If swallowed: | <ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If on skin or clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes and socks and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride (PVC).

Follow the manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply when weather conditions favor drift from treated areas. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding, or nectar-producing parts of plants if bees may forage on plants during this time period.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, shoes and socks, chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of United Turf Alliance, LLC. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, **THEN THIS PRODUCT SHOULD NOT BE APPLIED.** By applying this product Buyer acknowledges and accepts these inherent unintended risks **AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.**

United Turf Alliance, LLC shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

(continued)

(continued)

LIMITED WARRANTY

United Turf Alliance, LLC warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, UNITED TURF ALLIANCE, LLC MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of United Turf Alliance, LLC or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, United Turf Alliance, LLC or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF UNITED TURF ALLIANCE, LLC OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF UNITED TURF ALLIANCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements United Turf Alliance, LLC must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify United Turf Alliance, LLC of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

United Turf Alliance, LLC and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty** and **Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

RESISTANCE MANAGEMENT

ArmorTech® Guillotine™ contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **ArmorTech Guillotine** or other Group 4A.

To delay insecticide resistance consider:

- Avoiding the consecutive use of **ArmorTech Guillotine** or other Group 4A insecticides that have a similar target site of action on the same insect species.
- Using tank mixtures or premixes with insecticides from a different target site of action Group as long as the involved products are all registered for the same use and have different sites of action.
- Basing insecticide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitoring treated insect populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors and/or manufacturer for insecticide resistance management or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report a suspected resistance, you may contact United Turf Alliance, LLC at (770) 335-3015.

USE INFORMATION

ArmorTech Guillotine is a broad spectrum insecticide, for control of a wide spectrum of insects infesting turfgrass, ornamental plants, interior landscapes and non-bearing fruit and nut trees (in landscapes). When applied as directed, **ArmorTech Guillotine** provides excellent and long residual insect control.

RESTRICTIONS AND LIMITATIONS

- Do not use a foliar application of clothianidin (**ArmorTech Guillotine**) following a soil application of clothianidin.
- Regardless of formulation or type of application (soil or foliar), do not apply more than a total of 0.4 lb clothianidin per acre per season for turf, landscape, ornamentals and non-bearing fruits and nuts.
- If the maximum season limit of clothianidin as mentioned above has been applied and pest populations require additional treatments, use another registered pesticide that is not in the neonicotinoid class of chemistry.
- Do not apply by air.

PLANT TOLERANCE

Neither the manufacturer nor the seller has determined whether or not **ArmorTech Guillotine** can be used safely on cultivars of plants registered for use. **ArmorTech Guillotine** has been tested on many cultivars with no phytotoxicity observed at label rates. Since all plant species and their varieties and cultivars have not been tested for tolerance, it is recommended that a small number of plants be sprayed to make certain that no phytotoxicity occurs, prior to any large scale application to plants. The end user assumes all risks arising from application of **ArmorTech Guillotine** in a manner inconsistent with its labeling.

ArmorTech Guillotine can be tank mixed. However, all plant species and their varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments and adjuvants and surfactants. Conduct a spray mix compatibility and phytotoxicity trial under local conditions to ensure compatibility prior to any large scale use.

MIXING INSTRUCTIONS

1. Begin with clean equipment.
2. Add sufficient clean water to the spray tank for 1/2 of the mix load.
3. Start tank agitation.
4. Add and properly suspend the necessary amount of **ArmorTech Guillotine** before adding any other products. Agitate to ensure thorough mixing while adding the remaining required water. A high quality wetting agent or other spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions. Be sure to suspend and dilute **ArmorTech Guillotine** properly prior to the addition of any adjuvant. Consult adjuvant label or manufacturer for crop tolerance and safety information when used with this product.
5. Maintain agitation during mixing and application.
6. Apply with properly calibrated spray equipment.

SPRAY DRIFT MANAGEMENT

Do not allow this product to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions where states have more stringent regulations.

APPLICATION PROCEDURES

Foliar Application

Select spray nozzles which will provide accurate and uniform spray deposition. Use spray nozzles which provide medium sized droplets and reduce drift. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State and County Extension Service.

Apply **ArmorTech Guillotine** using sufficient water volume to provide thorough and uniform coverage. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. Do not make applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

TURFGRASS

ArmorTech Guillotine can be used as directed on grass grown in industrial, commercial and residential landscapes including airports, athletic fields, cemeteries, commercial sod farms, golf courses, home lawns, parks and playgrounds.

ArmorTech Guillotine has sufficient residual activity to provide high levels of control when applications are made proceeding or during the egg laying activity of the target pests. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Follow application with sufficient irrigation or rainfall to move the active ingredient through the thatch.

ArmorTech Guillotine can be applied to turf at 6.4 to 12.8 oz per acre. The rate is dependent on the target pest(s), their stage of development and the desired length of control. Multiple applications can be made but do not exceed the maximum amount per year (12.8 oz per acre). Consult your local State Agricultural Experiment Station, State Extension Turf Specialists or other turf experts for specific information concerning the timing of application(s). When targeting multiple pests, make application to earliest egg lay or adult flight. The high end of the rate range may be needed if extended control of secondary pests is desired.

Apply **ArmorTech Guillotine** in sufficient water volume (1 to 5 gallons per 1,000 sq ft) to provide optimal distribution in the treated area. The use of properly calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

APPLICATION ON TURFGRASS

PESTS	ARMORTECH GUILLOTINE APPLICATION RATES	TIMING INSTRUCTIONS
Annual Bluegrass Weevil (larval stages) Billbugs Black Turfgrass Ataenius <i>Phyllophaga</i> spp. (May or June Beetles) Spittle Bug White Grubs including: Asiatic Garden Beetle European Chafer Japanese Beetle Green June Beetle Northern Masked Chafer Oriental Beetle Southern Masked Chafer	6.4 to 12.8 oz/A (0.2 to 0.4 lb ai/A)	<p>Preventative Applications: Make applications through peak egg hatch of target species. Use the upper end of the rate range if application is made 60 days or more prior to peak adult flight and/or egg lay.</p> <p>Curative Applications: Curative applications are considered applications after egg hatch of the target species or when obvious turf damage from current generation has occurred. Use the upper end of the rate range when making applications post egg hatch.</p>
Armyworms Chinch Bugs including: Hairy Southern Crane Fly including: American European Cutworms Nuisance Ants (excluding Fire, Harvester, Carpenter, Pharaoh) Sod webworms	9.6 to 12.8 oz/A (0.3 to 0.4 lb ai/A)	<p>Crane Fly: Apply in the spring, when larvae are mature but prior to pupation or in the fall prior to egg hatch.</p> <p>Other pests: Apply when pest first appears.</p>
Mole Crickets (suppression)	12.8 oz/A (0.4 lb ai/A)	Time application to peak egg lay or early instars. Suppression can either mean control that is not commercially acceptable or inconsistent, ranging from poor to good.
<p>• Do not apply more than a total of 12.8 oz (0.4 lb ai) of ArmorTech Guillotine per acre per season.</p>		

ORNAMENTALS IN COMMERCIAL AND RESIDENTIAL OUTDOOR LANDSCAPE AND INTERIOR PLANTSCAPES AND OTHER SITES WHERE PLANTS ARE GROWN FOR AESTHETIC AND CLIMATE MODIFICATION PURPOSES

ArmorTech Guillotine is for use on ornamentals for insect control in commercial and residential landscapes and interior plantscapes. To assure optimum effectiveness, good spray coverage of the target plant is essential. Application can be made by foliar application or soil applications including soil injection, drenches, and broadcast sprays. For optimum control, make applications prior to anticipated pest infestation.

Apply **ArmorTech Guillotine** when threshold populations are observed. Do not wait until population beyond threshold has been established. Apply **ArmorTech Guillotine** in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve adequate coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.

Apply at planting or after planting to established plants. On established plants, irrigate the treated area thoroughly but do not allow leaching and runoff from containers for at least 7 days.

Choose lower rates for light infestations and/or smaller plants and higher rates for heavy infestations and/or larger plants.

RESTRICTIONS AND LIMITATIONS

- Do not apply more than a total of 12.8 oz (0.4 lb active ingredient clothianidin) of **ArmorTech Guillotine** per acre per season.
- Drench or broadcast applications will provide control of foliage feeding insects, however, control will not be achieved until the product has been taken up by the plant roots and translocated to the area where the insects are feeding.
- If making broadcast applications, water in with enough irrigation or rainfall to move the product into the zone where insects are active.
- For drench applications, use enough water to move the product into the zone where the insects are active.

**FOLIAR APPLICATION ON ORNAMENTALS IN
OUTDOOR LANDSCAPES AND INTERIOR PLANTSCAPES**

PLANTS	PESTS	ARMORTECH GUILLotine APPLICATION RATES (Per 100 Gals of Water)	SPECIAL INSTRUCTIONS
Bedding Plants Evergreens Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Ornamental Trees Shrubs	Aphid	0.63 to 1.26 oz	Uniformly apply the spray solution to the point of drip. Reapply as needed. The amount of spray solution needed per acre will depend on the ornamental size and the density of the leaf canopy.
	Mealybugs Whiteflies (suppression)	0.9 to 1.26 oz	
<ul style="list-style-type: none"> Do not exceed a total of 12.8 oz (0.4 lbs active ingredient clothianidin) of ArmorTech Guillotine per acre per season. 			

SOIL APPLICATIONS TO ORNAMENTALS IN OUTDOOR LANDSCAPES AND INTERIOR PLANTSCAPES

For systemic insect control on ornamental plants grown in-ground and outdoor landscapes (commercial, industrial, recreational and residential)

PLANTS	PESTS	ARMORTECH GUILLOTINE APPLICATION RATES	DIRECTIONS
Ornamental Plants including: Bedding Plants Evergreens Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Ornamental Trees Shrubs	Aphids Mealybugs Whiteflies (suppression) White Grubs including: Asiatic Garden Beetle European Chafer Green June Beetle Japanese Beetle Northern Masked Chafer Oriental Beetle Southern Masked Chafer	<p style="text-align: center;">Shrubs</p> 1.2 to 2.4 grams per foot of height 11.2 to 22.8 grams per 10 cumulative feet of height (0.38 to 0.78 oz per 10 cumulative feet of height)	<p>Soil Drench: Mix required dose in water and uniformly apply to soil around base of shrub or tree. Pull back mulch before drenching. For optimal performance, apply at least 2 quarts of dilute solution per foot of height. If lower drench volume is used, apply 1/2 inch of irrigation immediately after application to move product into root zone. Keep soil moist for at least 7 days.</p> <p>Soil Injection: Mix required dose in water and make at least four injections per shrub. For optimal performance, inject at least 2 qts of dilute solution per foot of height. Keep soil moist for at least 7 days.</p>
		<p style="text-align: center;">Multi-Trunk Trees</p> 1.2 to 4.8 grams per cumulative inch of trunk diameter at breast height (DBH) 11.2 to 48 grams per 10 cumulative inches of trunk diameter at breast height (DBH) (0.38 to 1.6 oz per 10 cumulative inches of trunk diameter at breast height (DBH)) <p style="text-align: center;">Single-Trunk Trees</p> 1.2 to 4.8 grams per inch of trunk diameter at breast height (DBH) 11.2 to 48 grams per 10 inches of trunk diameter at breast height (DBH) (0.38 to 1.6 oz per 10 inches of trunk diameter at breast height (DBH))	<p>Soil Drench: Mix required dose in water and uniformly apply to soil around base of tree, directed at the root zone. Pull back plastic or mulch before drenching. For optimal performance, apply at least 2 quarts of dilute solution per inch of trunk diameter. If lower drench volume is used, apply 1/2 inch of irrigation immediately after application to move product into root zone. Keep soil moist for at least 7 days.</p> <p>Soil Injection: Mix required dose in water and make at least four injections per tree with a low-pressure applicator. Use same amount of solution per hole. Injections can be made using the following methods: Grid System - Space injections on a 2.5 ft center extending to drip line. Circle System - Make injections in concentric circles extending inward from drip line. Basal System - Space injections evenly around trunk no more than 12 inches out from the base. Keep soil moist for at least 7 days after application.</p> <p style="text-align: right;">(continued)</p>

SOIL APPLICATIONS TO ORNAMENTALS IN OUTDOOR LANDSCAPES AND INTERIOR PLANTSCAPES

For systemic insect control on ornamental plants grown in-ground and outdoor landscapes (commercial, industrial, recreational and residential) (continued)

PLANTS	PESTS	ARMORTECH GUILLOTINE APPLICATION RATES	DIRECTIONS
Ornamental Plants including: Bedding Plants Evergreens Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Ornamental Trees Shrubs	Aphids Mealybugs Whiteflies (suppression) White Grubs including: Asiatic Garden Beetle European Chafer Green June Beetle Japanese Beetle Northern Masked Chafer Oriental Beetle Southern Masked Chafer	Broadcast Rates for Plants Grown in Beds: 4.3 to 8.3 grams per 1,000 sq ft (6.4 to 12.8 oz/A)	Make an application in such a manner as to maximize spray contact with soil. Use a sufficient water volume to incorporate the product into the soil, either during application or soon thereafter. Overhead irrigation immediately after application is recommended to move product off of plant foliage and to the soil surface.

ArmorTech Guillotine is a systemic insecticide and must move to the area of the plant where the target pest is feeding. Control of pests may be delayed following an application for one or more weeks as **ArmorTech Guillotine** moves to the feeding site. Factors such as plant size, soil type, and water demands will impact the time between application and obvious insect control.

- Do not apply more than 12.8 oz (0.4 lb ai) of **ArmorTech Guillotine** per acre per season.

ArmorTech Guillotine may be tank mixed or co-applied with other compatible insecticides. Please confirm product compatibility using the water volumes planned with a jar test prior to mixing up a large amount of material.

RESIDENTIAL APPLE AND PEAR TREES
(Not for use in commercial production sites)

Apply **ArmorTech Guillotine** when threshold populations are observed. Do not wait until population beyond threshold has been established. Apply **ArmorTech Guillotine** in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.

FOLIAR POST BLOOM APPLICATION RATES FOR USE ON RESIDENTIAL BEARING APPLE AND PEAR TREES

CROP	PESTS	ARMORTECH GUILLOTINE APPLICATION RATES	SPECIAL INSTRUCTIONS
Residential bearing: Apple Crabapple Pear	Aphids Leafhoppers Apple Maggot Leafminers Plum Curculio Codling Moth Leafrollers (suppression) Oriental Fruit Moth Soft Scales Pear Psylla	2 to 3 oz per 100 gallons of spray solution or 4 to 6 oz/A (2 to 3 oz ai/A) Rates per 100 gallons assume a spray rate of 200 gallons per acre. Adjust ArmorTech Guillotine rate per 100 gallons so as not to exceed 6 oz ArmorTech Guillotine per acre.	Apply ArmorTech Guillotine when threshold populations are observed. Do not wait until large insect populations have been established. The amount of ArmorTech Guillotine required per acre will depend on tree size and pest pressure. Choose lower rates for light infestation and/or small trees, and the higher rates for heavy infestations and/or larger trees. Apply by ground in 200 gallons per acre as a full coverage spray. For optimum performance of Leafminers and leafrollers, make the first application as soon as pol- lination is complete and bees are removed from the orchard. Repeat application as necessary.

- Do not apply treatments less than ten (10) days apart.
- Do not apply within seven (7) days of harvest.
- Do not apply more than 6.4 oz (0.2 lb ai) of **ArmorTech Guillotine** per acre per season.
- Do not feed or allow livestock to graze on cover crops from treated orchards.
- Applications must be made post bloom.

NON-BEARING FRUIT AND NUT TREES

(In landscapes which are not intended for use in commercial production)

Apply **ArmorTech Guillotine** when threshold populations are observed. Do not wait until population beyond threshold has been established. Apply **ArmorTech Guillotine** in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre.

APPLICATION RATES (POST-BLOOM APPLICATIONS) ON NON-BEARING FRUIT AND NUT TREES

PESTS	ARMORTECH GUILLOTINE APPLICATION RATES	SPECIAL INSTRUCTIONS
Aphids Apple Maggot Codling Moths Leafhoppers Plum Curculio Oblique-banded Leafrollers (suppression) Oriental Fruit Moth Spotted Tentiform Leafminers Western Tentiform Leafminers	2 to 3 oz per 100 gallons of spray solution or 4 to 6 oz/A (0.125 to 0.187 lb ai/A)	<p>Non-bearing fruit, including apples and pears, and nut trees in landscapes are only for ornamental use. FRUITS FROM THESE TREES ARE NOT TO BE USED AS FOOD AT ANY TIME.</p> <p>The amount of ArmorTech Guillotine per acre will depend on tree size, pest pressure and the volume of foliage present. The rate of water carrier per acre is based on a standard of 400 gallons of dilute spray per acre for large trees.</p> <p>For Control of Codling Moth and Oriental Fruit Moth: Apply ArmorTech Guillotine in a program of rotational sprays with other labeled insecticides. When using the 3 oz rate of ArmorTech Guillotine to control codling moth and/or oriental fruit moth, do not apply treatments less than a 14 day interval.</p> <p>Apply ArmorTech Guillotine in a full coverage spray in 200 gallons per acre to the point of drip. The spray volume will depend on tree size and density of the canopy. Repeat applications as needed to maintain control but do not apply more often than every 10 days.</p> <p>For Optimum Control of Leafminers and Leafrollers: Make the first application as soon as pollination is complete and bees are no longer present. Repeat application as necessary to maintain control. A single application may result in suppression only.</p>
<ul style="list-style-type: none"> • Do not apply more than 6.4 oz (0.2 lb ai) of ArmorTech Guillotine per acre per season. • Do not apply treatments less than ten (10) days apart. 		

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in a dry place away from excessive heat.

Do not store near food or feed.

Store in original container only.

To close package, replace and tighten cap to form an airtight seal.

Do not contaminate water, food or feed by storage or disposal.

For help with any spill, leak, fire or exposure involving this material, call day or night (800) 892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Copyright © 2010 by United Turf Alliance, LLC

ArmorTech is a registered trademark and Guillotine is a trademark of United Turf Alliance, LLC

Manufactured for

United Turf Alliance, LLC

8014 Cumming Highway

Suite 403-282

Canton, GA 30115

Form 1787-A Made in U.S.A.

EPA Reg. No. 59639-152-86064

EPA Est. 39578-TX-01[©], 67545-AZ-01[©]

Superscript is first letter in lot number.

ArmorTech®

GILLOTINE™

For foliar and systemic control of insects in commercial and residential landscapes and interiorscapes, non-bearing fruit and nut trees, residential apple and pear trees, turfgrass and other sites where plants are grown for ornamental, aesthetic and climate modification purposes

Active Ingredient	By Wt.
* Clothianidin	50.0%
Other Ingredients	50.0%
Total	100.0%
*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine	

EPA Reg. No. 59639-152-86064
EPA Est. 39578-TX-01(☉)
67545-AZ-01(☉) Superscript
is first letter in lot number.

**KEEP OUT OF
REACH OF
CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR
ADDITIONAL PRECAUTIONARY
STATEMENTS.

**NET WEIGHT
2 LB 8 OZ**

FIRST AID

- If in eyes:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If swallowed:**
- Call poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If on skin or clothing:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If inhaled:**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

Manufactured for **United Turf Alliance, LLC**
8014 Cumming Highway, Suite 403-282
Canton, GA 30115

Form 1787-A



United Turf Alliance