

Armor Tech

IMD 75

INSECTICIDE

**Insecticide for use in turfgrass and sod farms, and for landscape
ornamentals, and interior plantscapes.**

ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-*N*-nitro-2-imidazolidinimine . . . 75.0%

OTHER INGREDIENTS: 25.0%

TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT: 48 x 1.6 OZS.

EPA REG. NO. 228-588
EPA EST. NO. 67545-AZ-1

Manufactured for
Nufarm Americas Inc.
150 Harvester Drive
Burr Ridge, IL 60527

**To reorder contact
the United Turf Alliance
distributor
in your area.**

United Turf Alliance



FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

WPS USES: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses (e.g., crops, sod farms), must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Protective eyewear
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

NON-WPS USES: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, only agricultural plant uses are covered by the WPS, must wear:

- Shirt and pants
- Gloves
- Protective eyewear
- Shoes plus socks

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticide. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Protective eyewear
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

Keep children and pets off treated areas until dry.

RUNOFF MANAGEMENT

Do not cultivate within 10 feet of the aquatic areas to allow growth of vegetative filter strip. When used on erodible soils, use best management practices for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

ENDANGERED SPECIES NOTICE

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

RESISTANCE MANAGEMENT

Certain insects may develop resistance to insecticides after repeated use. Use different resistance management practices such as rotating classes of insecticides to help delay or minimize insect resistance.

This product contains the active ingredient imidacloprid, which is a Group 4A insecticide. Repeated use of Group 4A insecticides may lead to insects that become resistant to imidacloprid or other neonicotinoids (Group 4A) insecticides.

To reduce the chances of development of resistance to Group 4A insecticides, do not make more than three consecutive foliar applications of this product and/or other Group 4A insecticides with similar modes of action. In addition, Nufarm strongly recommends the use of other insecticides with a different mode of action prior to or after application of this

product. This strategy of insecticide rotation in concert with other IPM practices is considered an effective way to delay or minimize an insect's ability to develop resistance to this class of chemistry.

Some Group 4A neonicotinoid products used as foliar treatments include the active ingredients thiamethoxam (found in Actara® and Centric®), acetamiprid (found in Assail® and Intruder™), thiacloprid (found in Calypso®), and imidacloprid (found in Leverage®, Provado®, and Trimax™). Some 4A Group neonicotinoid products used as soil treatment include thiamethoxam (found in Platinum®) and imidacloprid (found in Admire®).

Additional information on insect resistance management may be obtained from your local extension specialist, certified crop advisor and/or product manufacturer, or from the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE [For product packaged in plastic containers]: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE STORAGE [For product packaged in Water-soluble Packaging]: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Exposure to moisture or excessive handling of water-soluble packets may cause breakage.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL [For product packaged in plastic container]: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container half full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONTAINER DISPOSAL [For product packaged in Water-soluble Packaging]: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Actara®, Centric®, and Platinum® are trademarks of a Syngenta Group Company.

Assail® is a trademark of Nippon Soda Co., Ltd.

Admire®, Merit®, Provado®, Calypso®, Leverage® and Trimax™ are trademarks of Bayer.

Bidrin® is a trademark of Amvac Chemical Corporation.

Intruder™ is a trademark of Nippon Soda Co., Ltd. and licensed for distribution by E.I. duPont de Nemours and Company.

Nufarm™ is a trademark of Nufarm.

ArmorTech IMD 75 Insecticide is not manufactured or distributed by Bayer CropScience, seller of Provado®.

ArmorTech IMD 75 Insecticide is not manufactured or distributed by Bayer Environmental Science, seller of Merit®.

PREPARATION OF SPRAY MIXES

This product is packaged in clear, water-soluble packets. Follow the instructions on handling these packets and for preparing spray solutions.

Handling Water Soluble Packets

The enclosed packets containing this product are water-soluble and will completely dissolve in water. To prevent the packets from dissolving before they are added to the spray tank, do not get the packets wet from handling with wet hands or wet gloves. The packets can break open if handled roughly. Remove the packets from the outer foil bag but do not open the packets. Keep unused packets sealed in the outer bag.

How to Prepare Spray Solutions

1. Fill the spray tank with 1/4 to 1/3 of the required amount of clear water and begin agitation.
2. Add the specified number of product packets. The packets should fully dissolve in 5 to 10 minutes. If water temperatures are cooler than normal, a longer amount of time may be required for the inner packets to completely dissolve.
3. Fill the tank with the remaining water needed. Maintain sufficient agitation during mixing and application.

If this product is to be tank-mixed with other pesticides and/or fertilizer solutions, check the compatibility (refer to the "Tank Mix Compatibility" section below) before adding to the spray tank. Use the following order of addition: 1) Product packets; 2) other wettable powders or wettable granules; 3) flowables (suspension concentrates); 4) emulsifiable concentrates. Run agitator as each component is added. Add the next component only after the previous one is thoroughly mixed. Then add the remaining amount of water to the spray tank. To ensure a uniform spray mixture, maintain constant agitation during both mixing and application.

Tank Mix Compatibility

This product has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. Before preparing tank mixtures with this product, especially if compatibility is not known, carry out the following small jar test using the desired tank mix partners.

1. Add proportionate amount of each component in the appropriate order to a pint or a quart jar.
2. Replace the cap, shake for 5 minutes, and allow the mixture to settle for 5 minutes.
3. Observe the jar for signs indicating an incompatible mixture. If the contents can be re-mixed by shaking and readily re-suspends, it is considered compatible. If the mixture separates out, foams, or forms a gel or lumps, then the mixture is not compatible.

PVA packets that are tank-mixed with products that contain boron or that release free chlorine will react to form a plastic that is insoluble in water or solvents (such as diesel oils, kerosene, gasoline or alcohol). **NOTE:** Chlorinated water may be used. Because the water-soluble packets are not soluble in petroleum-based liquids (such as kerosene, gasoline or alcohol), DO NOT attempt to use this product water-soluble packets directly in diesel oils or summer spray type oils such as those used in ULV or LV applications. Further information is available from your local Nufarm representative.

Restrictions

- DO NOT apply through any type of irrigation system.
- Keep children and pets off treated areas until dry.
- Do not allow livestock to graze in treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff of irrigation water. Do not allow puddling of irrigation water.
- **Rotation Crops:** Crops which are listed on imidacloprid labels or crops that have existing tolerances for imidacloprid may be planted in treated areas as soon as practical after the last imidacloprid application. Crop that are not found on an imidacloprid label, or crops that do not have existing tolerances for imidacloprid, may not be planted in treated areas for 12 months after the last application. Note that if cover crops are planted any time after an application of this product for soil building or erosion control those crops may not be grazed or harvested for food or feed.

Insect Resistance

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

TURF

This product will control or suppress insect pests that live in soil under lawns or grassy areas in residential and non-residential areas and sod farms (refer to table below for sites). Best control is obtained when applications are made before or during the egg-laying period. Irrigation is required after application to ensure residues of this product are moved through the thatch and into the soil layer. Refer to the table below for additional application instructions. Additional information on when to apply can be obtained from your local Agricultural Experiment Station, State Extension Turf Specialist, or Nufarm representative.

Sites	Area Treated With 1 Packet (1.6 oz.) of Product	Pests	Restrictions and Other Application Instructions
Turfgrasses around airports, athletic fields, cemeteries, golf courses, homes and multi-family residential buildings, office buildings or office parks, parks and playgrounds, shopping centers, and sod farms	8,250 - 11,000 sq. ft.	Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms (suppression only) European Chafer European Crane Fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	Application Instructions: Apply this product in sufficient water to ensure the turf receives an even uniform distribution of spray. Accurately calibrated equipment normally used for soil application of insecticides must be used and calibration must be checked often to ensure equipment works properly. Equipment that produces uniform, coarse droplet sprays with a low pressure setting will help to eliminate drift to non-target sites. In order for this product to be adequately distributed, do not apply the product to waterlogged grassy areas or to water-saturated soils. Rainfall or irrigation must occur within 24 hours of application to move this product vertically through the thatch and into the soil. Wait until after sufficient rainfall or irrigation has occurred to mow the grass. Annual Bluegrass Weevil, Billbugs, European Crane Fly, and Grubs: For best results, make applications before egg hatch.
	8,250 sq. ft.	Chinchbugs (suppression only) Mole Crickets	Chinchbugs: Make applications before hatching of first instar nymphs. Mole Crickets: Make applications before or during the peak egg-hatching period. This product may be applied with a curative insecticide when adults or large nymphs are present and actively tunneling. Restrictions: Do not apply more than 8.6 oz. (0.4 lb. of active ingredient) per acre per year.

TREES, ORNAMENTALS, GROUNDCOVERS AND INTERIOR PLANTSCAPES

This product may be applied by broadcast or foliar application to evergreens, flowers, foliage plants, groundcovers, interior plantscapes, non-bearing fruit and nut trees, ornamentals, shrubs, trees, vegetable plants intended for resale, and state, national, and private wooded forested areas (refer to table below for sites) to control or suppress insects. This product is a systemic insecticide that is absorbed by the roots and moves upward into the plant. For this product to control insects, it must come in contact with growing parts of the plant. Plant absorption of this product may be increased in some cases if it is applied with a fertilizer that contains nitrogen. Plants absorb this product from either foliar or soil applications. Refer to the table below for further instructions.

Woody Perennials: Protection in woody perennials is slower than in herbaceous species. Expect a delay of 2 or more weeks with longer delays for larger plants. Because of this, make applications to woody perennials well in advance of expected insect activity.

Bark Media: Treatments of this product to media with 30 to 50% or more bark content may confer a shorter period of protection.

Sites	Application Method and Number of Packets of Product (1.6 oz. Product) to Use	Pests	Restrictions and Other Instructions
Evergreens, Flowers, Foliage plants, Groundcovers, Interior plantscapes, Ornamentals, Shrubs, and Trees in and around residential, industrial, and commercial buildings and state, national, and private wooded and forested areas	<p>Foliar application: 1 packet in 300 gal. of water</p>	<p>Adelgids Aphids Japanese beetle (adult) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression only) Whiteflies</p>	<p>Application Instructions: Foliar Application: Apply this product in a sufficient volume of water to uniformly cover the treatment area. Foliar applications will provide systemic activity against target pests. If plants (such as holly, pine or ivy) have foliage that is difficult to wet, Nufarm recommends this product be applied with a spreader/sticker. Time applications to occur before heavy pest populations arise; make repeat applications as necessary.</p>
	<p>Broadcast application: 1 packet for 8,250 to 11,000 sq. ft.</p>	<p>White grub larvae (including Asiatic garden beetle, chafers, <i>Phyllophaga</i> spp., Japanese beetle larvae, and Oriental beetle)</p>	<p>Application Instructions: Broadcast Application: Mix the specified amount of this product in a sufficient volume of water to uniformly cover the treatment area. Apply in a minimum of 2 gallons of water per 1,000 sq. ft. After application, irrigate the treated areas to incorporate this product into the soil. Restrictions: Outdoor ornamentals: Do not apply by broadcast application more than 8.6 oz. (0.4 lb. active ingredient) per acre per year.</p>

Sites	Application Method and Number of Packets of Product (1.6 oz. Product) to Use	Pests	Restrictions and Other Instructions
<p>Trees in and around residential, industrial, and commercial buildings, Interior plantscapes, and state, national, and private wooded and forested areas</p>	<p>Soil Injection and Soil Drench: 1 packet per 24 to 48 inches of cumulative trunk diameter (DBH)</p>	<p>Adelgids Aphids Armored Scale (suppression only) Black vine weevil larvae Emerald Ash Borer Eucalyptus Longhorned Borers Flatheaded Borers (including bronze birch and alder borers) Japanese Beetles (adults) Lacebugs Leaf Beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips (suppression only) White grub larvae Whiteflies</p>	<p>Application Instructions: SOIL INJECTION: Use at least 4 holes per tree. GRID SYSTEM: Space injection holes on 2.5 ft. centers that extend to the drip line of the tree. CIRCLE SYSTEM: Evenly space injection holes in circles from the drip line in toward the trunk. More than one circle may be needed depending on the tree size. BASAL SYSTEM: Place injection holes evenly around the base of the tree trunk that extends only 6 to 12 inches from the base.</p> <p>Prepare this product in a sufficient volume of water so an equal amount of solution is injected into each hole using a low pressure. Use enough solution so that it reaches the root zone. Irrigation or rainfall for 7 to 10 days after application will provide optimum control.</p> <p>Restrictions: Do not apply this product by soil injection in Nassau or Suffolk Counties in the state of New York.</p> <p>Application Instructions: SOIL DRENCH: Before application, be sure there are no physical barriers (such as plastic tarp) present that may prevent the solution from reaching the root zone. Apply in a minimum of 10 gallons of water per 1,000 sq. ft. Apply the spray solution uniformly around the base of the tree ensuring the drench is directed at the root zone.</p> <p>Borers: If trees are heavily infested, an application of this product may not prevent the loss of the trees from existing pest damage and tree stress.</p> <p>Restrictions: Do not apply more than 8.6 oz. (0.4 lb. of active ingredient) per acre per year.</p>

(continued)

Sites	Application Method and Number of Packets of Product (1.6 oz. Product) to Use	Pests	Restrictions and Other Instructions
Shrubs in and around residential, industrial, and commercial buildings and state, national, and private wooded and forested areas	<p>Soil Injection and Soil Drench: 1 packet per 24 to 48 inches of cumulative shrub height)</p>		<p>Application Instructions: SOIL INJECTION: Use at least 4 holes per shrub. Prepare this product in a sufficient volume of water so an equal amount of solution is injected into each hole using a low pressure. Use enough solution so that it reaches the root zone of the individual shrubs. Irrigation or rainfall for 7 to 10 days after application will provide optimum control.</p> <p>Restrictions: Do not apply this product by soil injection in Nassau or Suffolk Counties in the state of New York.</p> <p>Application Instructions: SOIL DRENCH: Before application, be sure there are no physical barriers (such as plastic tarp) present that may prevent the solution from reaching the root zone. Apply in a minimum of 10 gallons of water per 1,000 sq. ft. Apply the spray solution uniformly around the base of the tree ensuring the drench is directed at the root zone.</p> <p>Restrictions: Do not apply more than 8.6 oz. (0.4 lb. of active ingredient) per acre per year.</p>

(continued)

Sites	Application Method and Number of Packets of Product (1.6 oz. Product) to Use	Pests	Restrictions and Other Instructions
Flowers and Ground covers in and around residential, industrial, and commercial buildings and state, national, and private wooded and forested areas	<p>Broadcast application: 1 packet for 8,250 to 11,000 sq. ft.</p>		<p>Application Instructions: Applications Prior to Planting Plants or to Established Plants: After application, this product must be incorporated into the soil. Irrigation to established plants after application will provide best results.</p> <p>Restrictions: Do not apply more than 8.6 oz. (0.4 lb. of active ingredient) per acre per year.</p>
To Manage Ants in the Ornamentals listed above	See above	Aphids Scale Mealy Bugs Other Sucking Insects	<p>Restrictions: When this product is used to control these target insects, ants are also controlled by limiting the honeydew available as a food source for the ants. This product may be used as a supplement to other commonly used methods (bait traps, residual sprays, etc.) that help eliminate unwanted ants in ornamentals.</p>

Crops	Use Rate and Pests	Restrictions and Other Instructions
<p>Pome Fruit in and around Residential Areas</p>	<p>1 Packet of product in 300 gallons of water (equivalent to 2.1 oz. product per acre)</p> <p>Aphids (except Woolly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs San Jose Scale</p>	<p>Pome fruit include: apple, crabapple, loquat, mayhaw, pear (including Oriental pear), quince</p> <p>Application Instructions: Foliar Application: Apply as needed after petal-fall. The use rate per acre is based on a standard of 400 gallons; therefore, adjust the amount of this product depending on the tree size and amount of foliage present.</p> <p>Rosy Apple Aphid: Apply prior to leaf rolling.</p> <p>Leafhopper: For late season (preharvest) control, apply while most leafhoppers are in the nymph stage.</p> <p>Leafminer: First generation – Apply as soon as pollination is complete and bees are removed from the orchard. For optimal control, apply as early as possible. Second and succeeding generations – Make applications early in the adult flight against egg and early instar larvae. For continued and severe pest pressure or overlapping generations, make a second application 10 days later. One application may only result in suppression. This product will not control late instar larvae.</p> <p>Mealybug: For best results be sure to thoroughly spray and cover the trunk and scaffolding limbs or other nesting sites.</p> <p>San Jose Scale: Time applications to the crawler stage and treat each generation. Do not use this product on this pest in pears in the state of California.</p> <p>Restrictions: California: Do not apply this product to control mealybugs and San Jose Scale on pears.</p> <p>Wait at least 10 days in between applications. Do not apply more than 2.1 oz. per A per application. Make only 5 applications per year. Harvest fruit 7 days or longer after the last application.</p>

Crops	Use Rate and Pests	Restrictions and Other Instructions
Pecans in and around Residential Areas	<p>1 Packet of product in 300 gallons of water (equivalent to 2.1 oz. product per acre)</p> <p>Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera</p>	<p>Application Instructions: Foliar Application: Apply as needed as pest pressure builds but before infestation is extremely heavy. The use rate per acre is based on a standard of 400 gallons; therefore, adjust the amount of this product depending on the tree size and amount of foliage present.</p> <p>Adequate control may be achieved only with two applications at 10- to 14-day intervals. For best results, a thorough and uniform spray coverage of foliage is necessary. To improve coverage, use an organosilicone-based spray adjuvant.</p> <p>Restrictions: Do not apply in the state of California unless otherwise directed by a supplemental label.</p> <p>Wait at least 10 days in between applications. Make only 3 applications per year. Do not apply more than 6.3 oz. of this product per acre per year.</p>
Ornamental Grapes in and around Industrial, and Commercial Buildings, and Residential areas	<p>Apply 1 Packet of product in 300 gallons of water (equivalent to 1.0 oz. product per acre)</p> <p>Leafhoppers (including glassy-winged sharpshooter) Mealybugs</p>	<p>Application Instructions: Apply as a foliar spray using 200 gallons of water per acre.</p> <p>Restrictions: Wait at least 14 days in between applications. Do not apply more than 2.0 oz. of this product per acre per year. Fruit may be harvested on the last day of application.</p>

ArmorTech IMD 75 INSECTICIDE

Insecticide for use in turfgrass and sod farms, and for landscape ornamentals, and interior plantscapes.

ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-*N*-nitro-2-imidazolidinimine 75.0%

OTHER INGREDIENTS: 25.0%

TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN No specific antidote is available. Treat the patient symptomatically.	

EPA REG. NO. 228-588
EPA EST. NO. 67545-AZ-1

Manufactured For
Nufarm Americas Inc.
150 Harvester Drive
Burr Ridge, IL 60527

NET WEIGHT: 48 x 1.6 OZS.

PULL HERE TO OPEN
PRESS TO RESEAL

630033

RV082908