



TEB 360 XL II

SPECIMEN LABEL

For control of listed diseases on ornamentals and golf courses

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethano..... 38.7%

OTHER INGREDIENTS..... 61.3%

TOTAL 100.0%

Contains 3.6 pounds tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to-mouth if possible. Call a poison control center or doctor for further treatment advice.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note To Physician: No specific antidote. Treat symptomatically. Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	

See inside booklet for additional PRECAUTIONARY STATEMENTS and complete DIRECTIONS FOR USE

EPA Reg. No. 42750-99

AD030121

SKU: 450711-1021



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170) The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pests out of treated areas until sprays have dried.

Chemigation: Do not apply this product through any type of irrigation system for turf and ornamental uses.

PRODUCT INFORMATION

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- Do not apply within 100 feet of aquatic areas listed above
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip
- See Spray Drift Management section for further information

SPRAY DRIFT MANAGEMENT

Make ground application when wind velocity favors on target product deposition (approximately 3 to 10 mph) Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume

For best results ArmorTech TEB 360 XL II may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label

Mixing

Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the ArmorTech TEB 360 XL II is completely dispersed in the mixture

Mixing Procedure

1. Water: Add three-quarters of the required volume to a thoroughly clean sprayer tank
2. Agitation: Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor: If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags: Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products: Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo emulsions (SE).
6. Water soluble products.
7. Emulsifiable concentrates (such as oil concentrates when applicable).
8. Water soluble additives (such as AMS or UAN when applicable).
9. Remaining quantity of water.

Resistance Management Information

The active ingredient in ArmorTech TEB 360 XL II is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and can exhibit cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development can not be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per year. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS

- Apply by ground application only in the state of New York.
- Do not make more than 3 applications at 0.6 fl. oz. of Tebuconazole 3.6FL (1.8 fl. oz.) (0.05 lbs AI) per 1000 sq. ft. per year in New York State. Do not apply more than 78 fl. oz. of Tebuconazole 3.6FL (2.2 lb) per acre per year in New York State.
- For use on golf course turf only
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (examples: elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 fl oz of ArmorTech TEB 360 XL II (0.1 lbs AI) per 1,000 sq ft per year.
- Do not apply more than 6 applications per year.

APPLICATION INSTRUCTIONS

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. ArmorTech TEB 360 XL II is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermuda grass can be sensitive to ArmorTech TEB 360 XL II under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid application when temperatures are expected to exceed 85 degrees F.

ArmorTech TEB 360 XL II can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of ArmorTech TEB 360 XL II in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, ArmorTech TEB 360 XL II should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

DISEASE	RATE (fl. oz./1000 Sq Ft)	NOTES
Dollar Spot <i>(Sclerotinia homoeocarpa)</i> Copper Spot <i>(Gloeocercospora sorghi)</i> Powdery Mildew <i>(Erysiphe graminis)</i> Corticium Red Thread <i>(Laetisaria fuaformis)</i> Rusts <i>(Puccinia spp.)</i> Red Thread <i>(Laetisaria fuaformis)</i> Pink Patch <i>(Limonomyces rosipellis)</i>	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. Do not make two consecutive applications of ArmorTech TEB 360 XL II or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Brown Ring Patch <i>(Waitea circinata)</i>	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development. Make two applications using the low to medium rate tank mixed with polyoxin D zinc salt fungicide at labeled rates. Repeat application at 21 day interval.
Brown Patch/Rhizoctonia Blight, Large Patch <i>(Rhizoctonia solani)</i> Anthracnose Basal and Foliar <i>(Colletotrichum cereale)</i>	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. Do not make two consecutive applications of ArmorTech TEB 360 XL II or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days
Bermuda Grass decline <i>(Gaeumannomyces graminis var. graminis)</i>	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Make two applications at the low to medium rate, 21 days apart or one application at the high rate. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide with a different mode of action as needed. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information.
Take All Patch <i>(Gaeumannomyces graminis var. avenae)</i>	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, make two applications at low rate, 21 days apart or one application at high rate in the fall when root zone soil temperatures reach 75° F and again in the spring at 55-60° F root zone soil temperatures. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

(continued)

DISEASE	RATE (fl. oz./1000 Sq Ft)	NOTES
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development (July Aug) using the low to medium rate. Do not make two consecutive applications of ArmorTech TEB 360 XL II or other DMI containing fungicides. Alternate with another fungicide with a different mode of action, such as thiophanate methyl. A second application may be made after 21 days.
Stripe Smut (<i>Ustilago striiformis</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	Make a single application to historical disease areas in spring as grass growth begins. For areas know to have higher disease pressure, use the medium to high rate.
Spring Dead Spot (<i>Leptosphaeria korrea</i> , <i>L narmari</i> , <i>Ophiosphaerella</i> <i>herpotricha</i> , <i>Gaeumannomyces</i> <i>graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrea</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, make two applications at the low to medium rate, 21 days apart or one application at the high rate in the fall when root zone soil temperatures reach 75° F. ArmorTech TEB 360 XL II can be combined with thiophanate-methyl at labeled rates. A spring application may be necessary in areas where disease pressure is known to be heavy. Apply using low to medium rate when root zone soil temperature reaches 55-60° F. Immediately after TEBUCONAZOLE 3.6FL is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	Apply first application in mid-June or 2-4 weeks prior time this blight normally becomes evident. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide with a different mode of action, such as thiophanate-methyl. A second application may be made after 21 days.
Summer Patch (<i>Magnaporthe poae</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention, apply in the spring when root zone soil temperatures reach 55-60° F. Do not make two consecutive applications of ArmorTech TEB 360 XL II or other DMI containing fungicides. Alternate with another fungicide with a different mode of action, such as thiophanate-methyl. Make additional applications at 21 day intervals. See local university recommendations for additional information. Immediately after fungicides is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the root zone.

(continued)

DISEASE	RATE (fl. oz./1000 Sq Ft)	NOTES
Zoysia Patch, Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	Make first application in early fall (mid September to mid October) prior to development of disease symptoms. Alternate with another fungicide with a different mode of action, such as polyoxin D zinc salt fungicide. Second and third applications may be made at 21 day intervals. See local university recommendations for additional information. An additional application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch (<i>Microdochium nivalis</i>)	0.6 – 1.1 (0.017 – 0.031 lb ai)	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that ArmorTech TEB 360 XL II be tank mixed with other registered snow mold products, such as those containing iprodione, thiophanate-methyl, polyoxin D zinc salt, or chlorothalonil, for best season long results.
Fairy Ring <i>Chlorophyllum (Lepiota), Lycoperdon, Marasmius</i> (Not For Use in California)	0.6 – 1.1 (0.017 – 0.031 lb ai)	For prevention in Cool season turf, make two applications at the low-medium rate in the spring when root zone soil temperatures reach 55-60° F. Make a second application using a 21-day interval. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Do not use a wetting agent unless hydrophobic soil conditions exist. For areas with historically high disease pressure, use the medium to high rate. For Warm season turf breaking dormancy, do not make two consecutive applications of ArmorTech TEB 360 XL II or other DMI containing fungicides. Alternate with other fungicides with a different mode of action. For hydrophobic areas, use an appropriate wetting agent to effectively penetrate the hydrophobic zone commonly created with this disease.
NOTE: Apply the specified amount of ArmorTech TEB 360 XL II in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of ArmorTech TEB 360 XL II per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS

- For use on ornamental plants only, not for woodlands or forest management
- For use only by professional applicators
- Do not apply more than 10 fl oz per acre in a single application
- Do not apply more than 0.31 gallons (40 fl oz) of ArmorTech TEB 360 XL II (1.13 lb AI) per acre per year
- Do not make more than 4 applications per year at highest rate
- Do not apply to bearing fruit trees or vegetables

ArmorTech TEB 360 XL II can be used in a disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when ArmorTech TEB 360 XL II is used in conjunction with sound disease management practices

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply ArmorTech TEB 360 XL II at rates of 4-10 fluid ounces (0.11 – 0.28 lbs AI) per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

NOTE: The “Directions for Use” of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

ORNAMENTALS DISEASE CONTROL

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14 - 21 days during the growing season, starting when leaves first appear	Apply every 14 days for a total of 3 applications beginning at the first sign of disease
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14 - 21 days apart, beginning with Spring bud break Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthracnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust		
Ground Covers and Vines	Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank
2. Close the sprayer, shake well and pressurize
3. Adjust nozzle to a coarse spray pattern and apply
4. Occasionally re pressurize the sprayer if needed to maintain a good spray pattern

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinseate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

In case of emergency call CHEMTREC at 1-800-424-9300.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Albaugh, LLC. All such risks shall be assumed by the user or buyer to the extent allowable under State law.

DISCLAIMER OF WARRANTIES: ALBAUGH, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Albaugh, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ALBAUGH, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.